arXiv:1606.07871v1 [math.NA] 25 Jun 2016

The Fourier expansion approximation for high-accuracy computation of the Voigt/complex error function at small imaginary argument

S. M. Abrarov* **and B. M. Quine***†

June 25, 2016

Abstract

It is known that the computation of the Voigt/complex error function is problematic for highly accurate and rapid computation at small imaginary argument $y << 1$, where $y = \mathrm{Im}[z]$. In this paper we consider an approximation based on the Fourier expansion that can be used to resolve effectively such a problem when $y \to 0$.

Keywords: Complex error function; Voigt function; Faddeeva function; complex probability function; spectral line broadening

The complex error function also known as the Faddeeva function is given by [1, 2, 3, 4, 5]

$$w(z) = e^{-z^2}\left(1 + \frac{2i}{\sqrt{\pi}}\int_0^z e^{t^2}dt\right), \tag{1}$$

where $z = x + iy$ is the complex argument. The real part of the complex error function is known as the Voigt function [1, 3, 6, 7] that are widely used in many fields of Physics, Chemistry and Astronomy. The integral on the

*Dept. Earth and Space Science and Engineering, York University, Toronto, Canada, M3J 1P3.
†Dept. Physics and Astronomy, York University, Toronto, Canada, M3J 1P3.

right side of equation (1) cannot be taken analytically. Consequently, the complex error function requires a numerical solution.

We have shown previously that using the Fourier expansion method the complex error function can be represented as [5, 8]

$$\begin{aligned} w\left(z\right) \approx & \frac{i}{2\sqrt{\pi}}\left[\sum_{n=0}^{N} a_n \tau_m \left(\frac{1-e^{i\ (n\pi+\tau_m z)}}{n\,\pi+\tau_m z} - \frac{1-e^{i\ (-n\pi+\tau_m z)}}{n\,\pi-\tau_m z}\right) - a_0 \frac{1-e^{i\tau_m z}}{z}\right] \\ = & i\frac{1-e^{i\tau_m z}}{\tau_m z} + i\frac{\tau_m^2 z}{\sqrt{\pi}} \sum_{n=1}^{N} a_n \frac{(-1)^n e^{i\tau_m z}-1}{n^2\,\pi^2-\tau_m^2 z^2}. \end{aligned} \tag{2}$$

where

$$a_n \approx \frac{2\sqrt{\pi}}{\tau_m} \exp\left(-\frac{n^2\pi^2}{\tau_m^2}\right)$$

is a set of the Fourier expansion coefficients, $\tau_m = 12$ and $N = 23$ (for simplicity we assume that $x = \mathrm{Re}\,[\mathrm{z}]$ and $y = \mathrm{Im}\,[\mathrm{z}]$ are non-negative).

The approximation (2) is efficient in computation. In particular, this approximation has been implemented in the latest version of the *RooFit* package for highly accurate and rapid computation of the Faddeeva function (C++ source code is provided in [9]). Furthermore, the equation (2) alone can cover with high-accuracy the entire domain $0 < x < 40,000$ and $10^{-4} < y < 10^2$ required for the HITRAN molecular spectroscopic database [10]. However, as the input parameter y decreases and becomes below 10^{-6}, the accuracy of the approximation (2) deteriorates. Although the deterioration in accuracy of approximation (2) with decreasing y is much slower as compared to the Weideman's approximation [2] (see also [5] for more information), it cannot cover the narrow band domain along x-axis at $y << 1$. It should be noted that the computation of the Voigt/complex error function at small $y << 1$ is a common problem in many approximations [6, 11, 12].

Let us show how a trivial rearrangement of the approximation (2) can effectively resolve this problem. Consider the following identity [1, 4]

$$w\left(z\right) = 2e^{-z^2} - w\left(-z\right)$$

that can be simply rewritten as

$$w\left(z\right) = e^{-z^2} + \frac{w\left(z\right) - w\left(-z\right)}{2}. \tag{3}$$

Substituting the approximation (2) into the right side of identity (3) immediately leads to

$$w\left(z\right)\approx e^{-z^{2}}-i\frac{\cos\left(\tau_{m}z\right)-1}{\tau_{m}z}+i\frac{\tau_{m}^{2}z}{\sqrt{\pi}}\sum_{n=1}^{N}a_{n}\frac{\left(-1\right)^{n}\cos\left(\tau_{m}z\right)-1}{n^{2}\pi^{2}-\tau_{m}^{2}z^{2}}. \qquad (4)$$

Theoretically, the equations (2) and (4) are identical. Practically, however, these equations cover different domains. In particular, while the approximation (2) provides high-accuracy at $y\gtrsim10^{-6}$, the approximation (4) is highly accurate at $y\lesssim0.1$.

In order to quantify accuracy of the approximation (4) we can define the relative errors for the real and imaginary parts as given by

$$\Delta_{\mathrm{Re}}=\left|\frac{\mathrm{Re}\left[w_{\mathrm{ref.}}\left(z\right)\right]-\mathrm{Re}\left[w\left(z\right)\right]}{\mathrm{Re}\left[w_{\mathrm{ref.}}\left(z\right)\right]}\right|$$

and

$$\Delta_{\mathrm{Im}}=\left|\frac{\mathrm{Im}\left[w_{\mathrm{ref.}}\left(z\right)\right]-\mathrm{Im}\left[w\left(z\right)\right]}{\mathrm{Im}\left[w_{\mathrm{ref.}}\left(z\right)\right]}\right|,$$

respectively, where $w_{\mathrm{ref.}}\left(z\right)$ is the reference. We obtained the highly accurate reference values by using Wolfram Mathematica (version 9) in enhanced precision mode.



Fig. 1. Logarithm of the relative error $\log_{10}\Delta_{\mathrm{Re}}$ for the real part of the complex error function at $y\leq0.1$.



Fig. 2. Logarithm of the relative error $\log_{10} \Delta_{\mathrm{Im}}$ for the imaginary part of the complex error function at $y \leq 0.1$.

Figure 1 shows the logarithm of the relative error for the real part of the complex error function. As we can see from this figure the worst accuracy is $\sim 10^{-13}$. The accuracy rapidly improves with decreasing x. Specifically, at $x \lesssim 2$ it becomes $\sim 10^{-16}$ as illustrated in Fig. 1 by dark-blue/purple area along y-axis.

Figure 2 depicts the logarithm of the relative error for the imaginary part of the complex error function. We can see from Fig. 2 that the accuracy is nearly perfect as the worst accuracy in the imaginary part is $\sim 10^{-15}$.

The computational test we performed shows that the approximation (4) provides high-accuracy within the domain $0 \leq x \leq 10$ and $10^{-14} \leq y \leq 10^{-1}$. Thus, combining together the equations (2) and (4) we have developed an algorithm that covers with high-accuracy the entire domain required for practical applications.

References

[1] A.K. Hui, B.H. Armstrong and A.A. Wray, Rapid computation of the Voigt and complex error functions, J. Quantit. Spectrosc. Radiat. Transfer, 19 (5) (1978) 509-516.
`http://dx.doi.org/10.1016/0022-4073(78)90019-5`

[2] J.A.C. Weideman, Computation of the complex error function, SIAM J. Numerical Analysis, 31 (1994) 1497-1518.
http://dx.doi.org/10.1137/0731077

[3] F. Schreier, The Voigt and complex error function: A comparison of computational methods. J. Quant. Spectrosc. Radiat. Transfer, 48 (1992) 743-762.
http://dx.doi.org/10.1016/0022-4073(92)90139-U

[4] M.R. Zaghloul and A.N. Ali, Algorithm 916: computing the Faddeyeva and Voigt functions, ACM Trans. Math. Software 38 (2011) 15:1-15:22.
http://dx.doi.org/10.1145/2049673.2049679

[5] S.M. Abrarov and B.M. Quine, Efficient algorithmic implementation of the Voigt/complex error function based on exponential series approximation, Appl. Math. Comp., 218 (2011) 1894-1902.
http://dx.doi.org/10.1016/j.amc.2011.06.072

[6] B.H. Armstrong, Spectrum line profilles: the Voigt function, J. Quant. Spectrosc. Radiat. Transfer. 7 (1967) 61-88.
http://dx.doi.org/10.1016/0022-4073(67)90057-X

[7] K.L. Letchworth and D.C. Benner, Rapid and accurate calculation of the Voigt function, J. Quant. Spectrosc. Radiat. Transfer, 107 (2007) 173-192.
http://dx.doi.org/10.1016/j.jqsrt.2007.01.052

[8] S.M. Abrarov and B.M. Quine, On the Fourier expansion method for highly accurate computation of the Voigt/complex error function in a rapid algorithm, arXiv:1205.1768 (2012).

[9] T.M. Karbach, G. Raven and M. Schiller, Decay time integrals in neutral meson mixing and their efficient evaluation, arXiv:1407.0748 (2014).

[10] L.S. Rothman, I.E. Gordon, Y. Babikov, A. Barbe, D.C. Benner, P.F. Bernath, M. Birk, L. Bizzocchi, V. Boudon, L.R. Brown, A. Campargue, K. Chance, E.A. Cohen, L.H. Coudert, V.M. Devi, B.J. Drouin, A. Fayt, J.-M. Flaud, R.R. Gamache, J.J. Harrison, J.-M. Hartmann, C. Hill, J.T. Hodges, D. Jacquemart, A. Jolly, J. Lamouroux, R.J. Le Roy, G.

Li, D.A. Long, O.M. Lyulin, C.J. Mackie, S.T. Massie, S. Mikhailenko, H.S.P. Müler, O.V. Naumenko, A.V. Nikitin, J. Orphal, V. Perevalov, A. Perrin, E.R. Polovtseva and C. Richard, The HITRAN2012 molecular spectroscopic database, J. Quant. Spectrosc. Radiat. Transfer, 130 (2013) 4-50.
http://dx.doi.org/10.1016/j.jqsrt.2013.07.002

[11] H. Amamou, B. Ferhat and A. Bois, Calculation of the Voigt function in the region of very small values of the parameter a where the calculation is notoriously difficult, Amer. J. Anal. Chem., 4 (2013) 725-731.
http://dx.doi.org/10.4236/ajac.2013.412087

[12] S.M. Abrarov and B.M. Quine, Accurate approximations for the complex error function with small imaginary argument, J. Math. Research, 7 (1) (2015) 44-53.
http://dx.doi.org/10.5539/jmr.v1n1p44